5

15

25

35

Claims

- 1) Live attenuated bacterium of the genus *Escherichia, Yersinia* or *Salmonella*, said bacterium not having a functional tRNA₅ leu, for use in a vaccine.
- 2) Live attenuated bacterium for use according to claim 1, said bacterium not having a functional tRNA₅ leu as a result of a mutation in the *leux* gene.
- 3) Live attenuated bacterium for use according to claim 1 or 2, wherein said bacterium is selected from the group consisting of *E. coli*, *S. enterica* serotype *typhimurium*, *enteritidis*, *choleraesuis*, *dublin*, *typhi*, *gallinarum*, *abortusovi*, *abortus-equi* or *pullorum*.
 - 4) Live attenuated bacterium for use according to claim 1-3, characterised in that the mutation comprises an insertion and/or a deletion.
 - 5) Live attenuated bacterium for use according to clams 1-4, characterised in that said bacterium carries a heterologous gene.
- 6) Live attenuated bacterium for use according to claim 5 characterised in that the heterologous gene is inserted in the *leux* gene.
 - 7) Live attenuated vaccine for the protection of animals and humans against infection with a pathogenic bacterium or the pathogenic effects thereof, characterised in that said vaccine comprises a bacterium as defined in claims 1-6 and a pharmaceutically acceptable carrier.
 - 8) Live attenuated vaccine according to claim 7, characterised in that it comprises an adjuvant.
- 9) Live attenuated vaccine according to claim 7 or 8, characterised in that it is in a freeze-30 dried form.
 - 10) Use of a live attenuated bacterium as defined in claims 1-6 for the manufacture of a vaccine for the protection of animals against infection with a pathogenic bacterium or the pathogenic effects of infection.

WO 2005/021031 PCT/US2004/027896

14

11) Method for the preparation of a vaccine according to claims 7-9, characterised in that said method comprises the admixing of a live attenuated bacterium as defined in claims 1-6 and a pharmaceutically acceptable carrier.